

What do we learn from Resonance Production in Heavy Ion Collisions ?

Christina Markert¹ † for the STAR collaboration

¹ Physics Department, Yale University,
New Haven, CT 06520, USA

Abstract. The resonance production of Δ , $K(892)$, $\Sigma(1385)$, $\Lambda(1520)$ and ϕ from elementary p+p and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV from the STAR experiment at RHIC is presented. Yields and spectra are discussed in terms of chemical and thermal freeze-out conditions. Thermal models do not describe sufficiently the yields of the resonance production in central Au+Au collisions. The approach which includes pseudo-elastic and elastic hadronic interactions after chemical freeze-out until thermal freeze-out suggests a time span of $\Delta\tau > 5$ fm/c.

Keywords: resonances, chemical freeze-out, thermal freeze-out, lifetime
PACS: specifications see, e.g. <http://www.aip.org/pacs/>

1. Introduction

Resonances have been proposed as a signature of a strongly interacting nuclear medium due to modification of their properties in the medium [1]. The leptonic decay products of resonances may provide a direct measurement of the hadronic stage due to their very small final state interaction cross section. However, hadronic decay products of resonances interact with hadrons of the medium. Therefore they maybe useful in describing the evolution of the source at this hadronic stage in terms of the temperature and lifetime between chemical and thermal freeze-out. The fact that in elementary p+p interactions the thermal freeze-out temperature, from a statistical model, is close to the chemical freeze-out temperature suggests that the life span is close to zero. Comparison of the resonance yields and spectra from Au+Au to p+p collision systems may indicate the influence of the hadronic interacting medium. Microscopic model calculations give suppression or enhancement factors for the resonance yields and modifications of the momentum spectra due to rescattering and regeneration of resonances in the hadronic medium [2, 3]. The suppression is present because the resonance invariant mass cannot be reconstructed by their decay daughter if one of them rescatters in the medium.

2. Data Analysis

The Δ , $K(892)$, $\Sigma(1385)$, $\Lambda(1520)$ and ϕ resonances are reconstructed by measuring their decay daughters ($\Delta \rightarrow p + \pi$, $K(892) \rightarrow K + \pi$, $\Lambda(1520) \rightarrow p + K$ and $\phi \rightarrow K + K$) with the STAR *Time Projection Chamber* (TPC). Charged decay particles are identified via energy loss (dE/dx) and their measured momenta. The neutral strong decays such as Λ from a $\Sigma(1385)$ decay are reconstructed via a topological analysis ($\Sigma(1385) \rightarrow \Lambda(\rightarrow p + \pi) + \pi$). The resonance signal is obtained by the invariant mass reconstruction of each daughter combination and subtraction of the combinatorial background calculated by mixed event or like-sign techniques [4]. The resonance ratios, spectra and yields are measured at mid-rapidity. The central trigger selection for Au+Au collisions takes 5% of the most central inelastic interactions. The setup for the p+p interactions is a minimum bias trigger.

3. Resonance yields in p+p and Au+Au collisions

The resonance and stable particle yields in p+p collisions at $\sqrt{s_{NN}} = 200$ GeV are well described by a statistical model with $T=170$ MeV [5]. This is expected if the particles are described by a statistical model in terms of phase space and if there is no hadronic phase after chemical freeze-out. The hadronic phase where the pseudo elastic and elastic interactions (rescattering and regeneration) occur dominantly. The statistical model fits the stable particles for $\sqrt{s_{NN}} = 200$ GeV Au+Au with $T=160-170$ MeV [6, 7, 8], but the resonances are not well described by the model predictions (see figure 1).

Microscopic model calculations (UrQMD [2, 3]) can describe the decrease for the $\Lambda(1520)/\Lambda$ (UrQMD: $\Lambda(1520)/\Lambda=0.03$) and the $K(892)/K^-$ (UrQMD: $K(892)/K^- = 0.25$), as they include rescattering of the decay daughters. UrQMD predicts a slight increase for the Δ^{++}/p from p+p to Au+Au collision systems. This is in agreement with the data (see figure 2), which means either the regeneration of the signal is of the same order as the rescattering or there is no rescattering. The Δ^{++} has a short lifetime of 1.3 fm/c, therefore the rescattering of the decay daughters is expected to be large. Due to the long lifetime of the ϕ resonance (44 fm/c), we would expect no significant signal loss by rescattering of the daughters, since most of the decays happen outside the fireball (10 fm/c), which is in agreement with our observation. The measured values of $K(892)/K$ and $\Lambda(1520)/\Lambda$ ratio suggest a lifetime interval between chemical and kinetic freeze-out greater than 5 fm/c when a thermal model with rescattering but no regeneration is used [9, 10].




Fig. 1. Particle ratios in central Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The lines are the predictions of a statistical model with $T = 160$ *pm* 5, $\mu_B = 24 \pm 4$, fitted to the stable particles for Au+Au collisions [6].

The decrease of the $\Lambda(1520)/\Lambda$ and $K(892)/K^-$ ratios from p+p to Au+Au collisions remains nearly constant from peripheral collisions (50-80%) up to the most central collisions (5%) (see figure 2). This result suggests the same $\Delta\tau$ for peripheral and central Au+Au collisions the same lifetime of the system between chemical and thermal freeze-out.




Fig. 2. Resonance/non-resonance ratios of ϕ/K^- [11], Δ^{++}/p [12], ρ/π [13], $K(892)/K^-$ [14] and $\Lambda(1520)/\Lambda$ [4, 15] for p+p and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The ratios are normalized to the ratio of $K^*(892)/K^-$ in p+p collisions. Statistical and systematic errors are included.

4. Resonance momentum spectra in p+p and Au+Au collisions

The mean transverse momentum, $\langle p_T \rangle$, as a function of mass in figure 3 [16] (left panel) shows a mass dependence for stable (circles) and resonance (squares) particles. A fit to the ISR data shown by the curve includes π , K and p from p+p collisions at $\sqrt{s_{NN}} = 26$ GeV [17] and is in good agreement with the π , K and p data from $\sqrt{s_{NN}} = 200$ GeV. The resonances with masses higher than 1 GeV/ c^2 and the Ξ indicate a stronger $\langle p_T \rangle$ dependence which is not represented by the fit to the ISR data. The higher mass particles are more dominantly produced in the higher multiplicity events which introduces an increase in their $\langle p_T \rangle$ and a bias in the event selection.

According to microscopic model predictions the loss of signal in the low momentum region would result in an increase of the inverse slope parameter (and $\langle p_T \rangle$) of the transverse momentum spectra for the resonances. The $\langle p_T \rangle$ of the stable particles shows a smooth rise from p+p to the most central Au+Au collisions. while the resonances show a steep rise and stay constant up to the most central Au+Au collision. This observation is in agreement with the rescattering and regeneration of the resonance signal in the low momentum region. The effect is much more pronounced for the $K(892)$ than for the ϕ , due to the much shorter lifetime of $K(892)$ and the

larger probability of signal loss due to rescattering. This is also in agreement with the measured yields of the resonances.




Fig. 3. Left: Mean transverse momentum, $\langle p_T \rangle$, as a function of particle mass in p+p collisions at $\sqrt{s_{NN}} = 200$ GeV. The curve is a fit to measured ISR p+p data (π , K and p) at a collision energy of $\sqrt{s_{NN}} = 26$ GeV [17]. Right: Mean transverse momentum $\langle p_T \rangle$ as a function of charged particles for p+p and Au+Au collision systems for π , K, p, ϕ , K(892) and Δ .

5. Conclusions

Resonances with their hadronic decay daughters are a unique tool to probe the time span, $\Delta\tau$, between chemical and thermal freeze-out in heavy ion collisions. The measured $\Lambda(1520)/\Lambda$ and $K(892)/K$ together with a thermal model and rescattering suggest a lower limit of $\Delta\tau > 5$ fm/c. The effect of rescattering in the transverse momentum distribution of the resonances can be observed as an increase of their $\langle p_T \rangle$ from p+p to most peripheral Au+Au collisions. The mass dependence of $\langle p_T \rangle$ in p+p collisions including the high mass particles indicates a linear trend, which is different from the ISR parameterization.

Acknowledgement(s)

At first I would like to thank the organizers who invited me to such a well organized workshop at such a nice place. My special thanks go to my theory friends Jörg Aichelin, Marcus Bleicher and Horst Stöcker. I would also like to thank Olga Barannikova, Sevil Salur and Rene Bellwied for the many good and exiting discussions that we had. And many thanks goes to the STAR collaboration for the

support in presenting this data.

References

1. R. Rapp and J. Wambach, *Adv. Nucl. Phys.* **25**, 1 (2000).
2. M. Bleicher and J. Aichelin, *Phys. Lett.* **B530** (2002) 81, hep-ph/0201123.
3. M. Bleicher and H. Stöcker, *J.Phys.* **G30** (2004) 111, private communication.
4. L. Gaudichet et al., *J.Phys.* **G30** (2004) 549.
5. F. Becattini, *Nucl. Phys.* **A702**, 336 (2002).
6. O. Barannikova, this proceedings.
7. P. Braun-Munzinger, D.Magestro, K. Redlich and J.Stachel, *Phys. Lett.* **B518** (2001) 41.
8. P. Braun-Munzinger et al., nucl-th/0304013.
9. G. Torrieri and J. Rafelski, *Phys. Lett.* **B509** (2001) 239-245.
10. C. Markert, G. Torrieri, J. Rafelski, Campos do Jordao 2002, New states of matter in hadronic interactions 533-552, hep-ph/0206260.
11. J. Ma et al., *J.Phys.* **G30** (2004) 543.
12. H. Zhang et al., nucl-ex/0403010.
13. P. Fachini et al., *J.Phys.* **G30** (2004) 565.
14. H. Zhang et al., *J.Phys.* **G30** (2004) 577.
15. C. Markert et al., Proc. 19th Winter Workshop on Nuclear Dynamics (2003) 71.
16. C. Markert et al., Quark Matter 2004, Oakland, California, 11-17 Jan 2004. nucl-ex/0404003.
17. M. Bourquin and J.-M. Gaillard, *Nucl. Phys.* **B114** (1976) 334.